

Trends in Fish Records

Margaret Ashton

TICK TOP

Douglas Herdson

Marine Fish Information Services *Douglas.Herdson@btinternet.com* South West Marine Ecosystems 2019, Plymouth, 12th April 2019





Britain's First Jellynose



1.9 m of Loppe's Tadpole fish *Ijimaia loppei* Plymouth Fish Market, April 2013







Pelvic fins

Pompano Dolphinfish

Mark Padfield

Coryphaena equiselis, Chesil beach 8 Sept 2017,



Crocodile shark *Shark Pseudocarcharias kamoharai*, Hope Cove, Devon. 18 February 2017



Crocodile shark First European report

Smalltooth Sand Tiger or ragged-tooth (*Odontaspis ferox*), Normandie. 13 August 2012

E Stephan

211

- E Stephan

Range extension



Red Mullet Mullus surmuletus



Juvenile Red Mullet Plymouth Sound Bioblitz, September 2014



Almaco Jack Seriola rivoliana



Blue Runner Caranx crysos



Derbio, *Trachinotus ovatus* Probably the seventh recorded in Britain

Plymouth Fish Market, 14th December 2018 Four different species of sea bream (Sparidae). Not a large amount of any, but this is the first time I can remember this occurring. There were at least six Couch's Pagrus pagrus, three or four Black Sea Bream Spondyliosoma cantharus, four nice Giltheads Sparus aurata, and one Red or Blackspot Pagellus bogaraveo.



Red (above) and Couch's (below) Sea Breams



Gilthead Bream Sparus aurata



Black Sea Bream Spondyliosoma cantharus



Pandora *Pagellus erythrinus* Whitsand Bay, September 2018.

Triggerfish, Forth Towan/ 2007

Emily Priestley



Atlantic bonito Sarda sarda



Figure 6. Specimen of Atlantic bonito (Sarda sarda) collected during the survey.

Bonitos had not previously been seen, but 91 0-year group (15 – 30 cm) were caught during the CEFAS PELTIC 18 cruise October-November 2018



Haddock Melanogrammus aeglefinus

Non-Native Fish



Rainbow trout, Onchorhynchus mykiss




Humpback Salmon, *Oncorhynchus gorbuscha*, Mid Norway, Pacific Pink or Humpback Salmon, Oncorhynchus gorbuscha caught in the River Camel, Cornwall September 2007

[Also north east Scotland, Northumberland and Co Durham]



Green Sunfish, Pumpkinseed or Rock Bass ? Mevagissey Harbour, Cornwall, 21 October 2018 Rock Bass Ambloplites rupestris



But some unexpected stories can emerge

Anchovy goldrush: UK trawlers cash in as warm water fish arrive by the ton

Daily Mail 31 October 2009





Two Devon trawlers hauled in a staggering 36 tons of anchovies - worth £72,000 - in just two days

The Guardian Tuesday 8 May 2012

Warm water marine species spreading northwards into British waters Higher sea temperatures around UK mean bluefin tuna,




thresher sharks and anchovies among others are increasingly common.

Exotic warm water marine species such as **anchovy**, bluefin tuna, stingray, and

thresher shark are spreading northwards into British coastal waters, where average sea temperatures are now moving closer to the warmer conditions of southern Europe. Severin Carrell, Scotland correspondent

Anchovy Engraulis encrasicolus



First described from Cornwall by Jonathan Couch in 1838.



November 1889, Large catches of sprats off Dover and Deal, with 25% of the catch being anchovies



Dover Beach, 1890

Anchovies in the English Channel By J T Cunningham, M.A.,F.R.S.E. Journal of the Marine Biological Association of the UK 1(03):328 -339; 1890

The Migration of the Anchovy By J T Cunningham, M.A. Journal of the Marine Biological Association of the UK 3(04):300 -303; 1895 Cunningham's studies suggested that the anchovies on the south coast in 1889 and 1890 had originated from the spawning stocks in the Zuider Zee, where they spawned around June and July.







Anchovy eggs

The main spawning area was lost in 1932 when the Zuider Zee was closed by a dyke to give polders. This left a remnant breeding population in the adjacent Wadden Sea.

North Sea catches of anchovy the remained at a minimal level until the 1990s; when they began to increase both there and in the Baltic, giving welcome catches to fishermen on the English east coast and even in July 2014 to anglers in Essex.



Juvenile Anchovy 44mm, Southampton water, 5 October 2016



PELTIC 2014 30th September –19th of October 2014

Unloading Anchovies







Anchovies *Engraulis encrasicolus* Plymouth, November 2014



Anchovies, Mounts Bay, 18 January 2017

Hence in a typical year anchovies occur off Portland in October. With the shoals off Plymouth in November/December and in Cornish inshore waters in December/January.

Two studies published in 2012 looked at single nucleotide polymorphisms and found that the North Sea and English Channel shoals were genetically homogeneous. Three Biscay stocks were also homogeneous; but there were significant genetic differences between the Biscay and North Sea and Channel stocks.




Figure 5. Acoustically derived anchovy biomass distribution (left) and trawl-based length frequency histogram for anchovy in each of the subareas of the Peltic survey. Please note that bubble size has not been standardised between species.

PELTIC 2018 (October/November 2018) found anchovies more widespread than previous years, with small fish in French waters. Not a typical year.

Many thanks for information from:-

Jeroen Van Der Kooij Peter Howlett John Lancaster Adam Powell Lawrence Raybone/Ilfracombe Aquarium **Paul Gainey** Nigel Jones Environmental Record Centre for Cornwall and the Isles of Scilly MarLIN **Oliver Crimmen** Marine and Fisheries Agency (now the Marine Management **Organisation**)

And many others



